



## Tennis specific limitations in players with an ACL deficient knee

J Maquirriain and P J Megey

*Br. J. Sports Med.* 2006;40:451-453  
doi:10.1136/bjasm.2005.023390

---

Updated information and services can be found at:  
<http://bjsm.bmjournals.com/cgi/content/full/40/5/451>

---

*These include:*

### References

This article cites 14 articles, 3 of which can be accessed free at:  
<http://bjsm.bmjournals.com/cgi/content/full/40/5/451#BIBL>

### Rapid responses

You can respond to this article at:  
<http://bjsm.bmjournals.com/cgi/eletter-submit/40/5/451>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Topic collections

Articles on similar topics can be found in the following collections  
[Sports Medicine](#) (1226 articles)

---

### Notes

---

To order reprints of this article go to:  
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *British Journal of Sports Medicine* go to:  
<http://www.bmjournals.com/subscriptions/>

## ORIGINAL ARTICLE

## Tennis specific limitations in players with an ACL deficient knee

J Maquirriain, P J Megey

*Br J Sports Med* 2006;40:451–453. doi: 10.1136/bjism.2005.023390

**Background:** Complete rupture of the anterior cruciate ligament (ACL) causes significant alteration of knee joint kinematics. Untreated patients often develop joint instability, chronic articular degeneration, and knee dysfunction. Demands on the ACL produced by playing tennis have not been investigated.

**Objective:** To identify subjective sport-specific limitations in tennis players with isolated unilateral ACL deficiency.

**Study design:** Prospective case-control study.

**Methods:** 16 players (mean (SD) age, 39.9 (2.3) years; 14 men) with a chronic unilateral ACL deficient knee and 16 healthy controls (38.25 (8.47) years; 14 men) were recruited. ACL deficiency was confirmed by clinical and magnetic resonance imaging. A Lysholm score was obtained in all patients, together with subjective evaluation of their current tennis performance compared with pre-injury levels, applying a 0–100% visual scale. Both groups completed a questionnaire on tennis specific abilities.

**Results:** Lysholm scores were: 85.6 (10.3) points in the study group and 100 (0) points in the control group ( $p < 0.001$ , *t* test for independent samples). Injured players evaluated their current tennis performance as 66.8 (15.2)% compared with 100% pre-injury level ( $p < 0.005$ , *t* test for dependent samples). Abilities affected in the ACL deficient group were landing after a smash stroke ( $p < 0.001$ ); stopping abruptly and changing ( $p < 0.001$ ); playing a three set singles match ( $p < 0.05$ ); and playing on a hard court surface ( $p < 0.001$ , Kolmogorov-Smirnov test).

**Conclusions:** There are specific limitations associated with complete isolated ACL rupture, including subjective tennis performance impairment, limitations landing after a smash, stopping and changing step direction, difficulties playing a three set singles match, and playing on hard court surfaces.

See end of article for authors' affiliations

Correspondence to:  
Dr Javier Maquirriain,  
National High  
Performance Training  
Centre (CeNARD),  
Argentine Tennis  
Association, Pilar, Buenos  
Aires, Argentina;  
[jmaquirriain@yahoo.com](mailto:jmaquirriain@yahoo.com)

Accepted 7 December 2005

Of all the ligaments of the knee, the anterior cruciate (ACL) is the one that most commonly suffers complete disruption. Untreated, this injury can cause potentially debilitating consequences including meniscus tears, anterior and rotational instability, and early onset osteoarthritis.<sup>1,2</sup> Consequently, most orthopaedic surgeons advocate surgical reconstruction of the ACL for patients seeking to take part in activities that exert high functional demands on this ligament. However, demands made on the ACL from tennis playing remain unknown, and ACL insufficiency in tennis players has generated little research interest, with only limited references in the sports medicine literature.

ACL rupture is not a common injury during tennis playing. Kühne *et al*<sup>3</sup> have reported a 1.8% overall ACL rupture incidence from tennis related injuries. Nevertheless, many people play tennis after suffering cruciate ligament rupture during other activities, mainly contact sports. These patients often seek medical advice about potential limitations of practising tennis with an ACL deficient knee.

Our aim in this study was to identify specific subjective limitations in tennis players with unilateral ACL deficiency.

## METHODS

Sixteen tennis players (mean (SD) age, 39.93 (2.33) years; two women, 14 men) with a unilateral ACL deficient knee and 16 healthy players as controls (mean age 38.25 (8.47) years; two women, 14 men) were analysed in a prospective study. Inclusion criteria for the patients were: age 18 years or older; unilateral isolated ACL deficient knee confirmed by clinical and magnetic resonance imaging (MRI); regular tennis practice (at least once a week) at any level; no previous lower limb surgery or major trauma; and ACL rupture at least

six months prior to the study period. For the controls the inclusion criteria were: age 18 or older; tennis practice on a regular basis at any level; no prior lower limb surgery or major trauma; and stable knees. Subjects in the control group were matched with the study group for age, sex, and tennis participation.

The clinical diagnosis of ACL deficiency was supported both by history and by physical examination, using commonly accepted anterior knee instability tests (Lachman test, pivot-shift tests, anterior drawer test). Physical examination revealed no evidence of additional knee joint instability (that is, medial or posterolateral instability).

MRI evidence of ACL rupture was based on primary and secondary signs following the guidelines of Vahey *et al*.<sup>4</sup>

Knee function evaluated by modified Lysholm score<sup>5</sup> generated numbers from 0–100 (highest) based on a series of questions pertaining to knee function. All 16 patients were asked to evaluate their tennis performance subjectively in relation to their preinjury level on a 0–100% visual scale.

Both groups responded a questionnaire on tennis specific abilities, adapted from Mont *et al*.<sup>6</sup> Each ability was evaluated using a scale of 1 to 5: 1, “cannot do it”; 2, “extremely difficult” (pain and instability); 3, “difficult” (instability only and slight pain); 4, “able to do it with some instability and no pain”; 5, “able to do it with no pain and no instability”. Any playing technique changes generated because of mistrust, fear of knee giving way, or other joint related complaints were graded as 1.

For simplification, the knee on the same side as the arm used for the serve was considered “dominant”. For statistical significance,  $\alpha$  error was calculated as 0.05 and  $\beta$  error as

**Abbreviation:** ACL, anterior cruciate ligament

0.20. Power analysis was estimated using software.<sup>7</sup> The Kolmogorov-Smirnov test for non-parametric variables of small independent samples was used for tennis abilities analysis. A standard statistical package (Statistics for Windows, StatSoft Inc, Tulsa, Oklahoma, USA) was used to generate all statistics.

## RESULTS

The right knee was affected in five players (31%) and the left knee in 11 (69%); five knees were dominant and 11 were non-dominant. ACL injury was caused by soccer (n = 10), skiing (n = 2), tennis (n = 1), paddle tennis (n = 1), rugby (n = 1), and others (n = 1).

Lysholm scores were 85.6 (10.3) points in the study group and 100 (0) points in the control group ( $p < 0.001$ , *t* test for independent samples).

Injured players evaluated their tennis performance as 66.9 (15.3)% compared with 100% preinjury levels ( $p = 0.0014$ , *t* test for dependent samples).

Abilities significantly affected in the ACL deficient group were: landing after a smash stroke ( $p < 0.001$ ); stopping abruptly and changing direction ( $p < 0.001$ ); playing a three set singles match ( $p < 0.05$ ); and playing on hard courts ( $p < 0.001$ ). No other abilities investigated showed statistically significant differences. All tennis ability assessments are shown in table 1.

No significant relation between limitations in baseline side movements (right, left) and the ACL deficient knee was found in this group of tennis players.

Time elapsed between ACL injury and return to tennis averaged 6.4 (9.8) months; time elapsed between ACL injury and study assessment averaged 40.2 (47.0) months. Only one

patient used a rigid knee brace for playing tennis. All players played at recreational level.

## DISCUSSION

The results of this study identified subjective limitations of subjects with ACL deficient knees during tennis movements. Tennis players did not admit to significant impairment when performing the majority of tennis strokes such as forehand, backhand, volleys, and serves. The major limitation referred to was landing after hitting a smash. This task has often been perceived by ACL deficient subjects as being difficult to carry out with confidence.<sup>8</sup> Many players from the study group referred avoiding the smash entirely.

Interesting information was obtained about displacement while playing tennis. Injured players did not have significant impairment in forward running such as in "serve and volley" or "reaching a drop shot" movements. This finding was in agreement with previous studies—for example, Czerniecki *et al* found that running in a straight line may not generate sufficient rotational torque to initiate rotational instability in the cruciate deficient knee.<sup>9</sup> In contrast, ACL deficient players show major limitation when trying to "stop suddenly and change direction". This type of stressful deceleration creates high anterior loading on the tibiofemoral joint.<sup>10</sup> Both external varus-valgus and internal-external rotation place increased load on the knee joint during cutting movements compared with normal running.<sup>11</sup> Varus-valgus and internal-external rotational movements are believed to be responsible for increasing knee joint ligament risk of injury. External flexion loads, valgus and internal rotation during side-stepping all have the potential to increase ACL and medial collateral ligament load substantially.<sup>11</sup> The ability of normal subjects to undertake deceleration tasks without ACL rupture or giving way of the knee is attributed to the coordinated interactions among the ligamentous and other soft tissue passive restraints, joint geometry and congruency, friction between cartilage surfaces, active muscular control, and tibiofemoral joint compressive forces.<sup>10</sup> Approximately 86% of shear forces are considered to be restrained by the ACL<sup>12</sup>; however, in ACL deficient knees these loads must be restrained by the articulating surfaces and the surrounding soft tissues.<sup>10</sup>

Most ACL injuries are indirect in nature yet occur during contact sports. Tennis involves tremendous forces during cutting, pivoting, and sudden deceleration manoeuvres; nevertheless ACL injuries are less common in tennis than in contact sports.<sup>13 14</sup> In this series, most injuries occurred during contact sports such as soccer or rugby, but the exact mechanism (direct *v* indirect) was not determined. Sallay *et al*<sup>13</sup> hypothesised that a tennis player is not as likely to sustain an ACL injury because of the ability of the neuromuscular system to coordinate muscular function in anticipation of each movement, with little surprise effect. Many investigators have indicated that anticipating a movement can change reflex responses and postural adjustments to minimise forthcoming perturbation and maintain appropriate posture. Besier *et al*<sup>15 16</sup> were able to confirm previous hypotheses indicating that knee joint moments increase under unanticipated conditions compared with preplanned manoeuvres, primarily because of a large increase in varus-valgus and internal-external rotational moments under unanticipated conditions. It is believed that unanticipated movement alters the external moments applied to the knee by reducing the time to implement appropriate postural adjustment strategies. Tennis may cause a low incidence of indirect ACL injury owing to the absence of frequent complete twisting manoeuvres and high jumping, as well as enough time for the player to anticipate strokes, especially from the baseline.

**Table 1** Subjective evaluation of tennis abilities (adapted from Mont *et al*<sup>6</sup>)

Ability	ACL deficient knee*	Control	p Value†
Bending to hit a shot at knee level	4.81 (0.75)	5.00 (0)	$p > 0.10$
Shifting weight in forehand	4.43 (1.03)	5.00 (0)	$p > 0.10$
Shifting weight in backhand	4.43 (0.89)	5.00 (0)	$p > 0.10$
Shifting weight in first serve	4.62 (0.71)	5.00 (0)	$p > 0.10$
Shifting weight in second serve	4.75 (0.68)	5.00 (0)	$p > 0.10$
Shifting weight in volleys	4.68 (0.79)	5.00 (0)	$p > 0.10$
Shifting weight in smash	2.68 (1.66)	5.00 (0)	$p < 0.001$
Moving to right side on the baseline	4.43 (0.81)	5.00 (0)	$p > 0.10$
Moving to left side on the baseline	4.37 (0.88)	5.00 (0)	$p > 0.10$
Moving forward to reach a drop-shot	4.50 (1.09)	5.00 (0)	$p > 0.10$
Moving forward after serve to volley	4.75 (0.44)	5.00 (0)	$p > 0.10$
Stopping abruptly and turning direction	3.50 (1.15)	5.00 (0)	$p < 0.001$
Hitting a ground stroke on the run	4.68 (0.70)	5.00 (0)	$p > 0.10$
Playing a 3-set match of singles	3.93 (1.38)	5.00 (0)	$p < 0.05$
Playing a 3-set match of doubles	4.68 (1.01)	5.00 (0)	$p > 0.10$
Playing on hard court	3.62 (1.54)	5.00 (0)	$p < 0.005$
Playing on clay court	4.81 (0.50)	5.00 (0)	$p > 0.10$

\*Graded scale: 1, cannot do it; 2, extremely difficult (pain and instability); 3, difficult (instability only and slight pain); 4, able to do it with some instability and no pain; 5, able to do it with no pain and no instability.  
†p values obtained by Kolmogorov-Smirnov test for non-parametric variables of small independent samples.

ACL injured players described significant impairment of their recreational tennis performance compared to preinjury level. Results from the present study may support the need for surgical treatment for competitive tennis players with ACL deficiency. Further studies are needed to determine the true incidence of ACL injuries in tennis, to analyse tennis motion knee biomechanics, and to establish the degree of improvement in tennis ability after ACL reconstruction.

Tennis players with an ACL deficient knee showed a clear incapacity to play on hard courts, where demanding eccentric deceleration motions occur. Frontal and rotational knee moments are thought to be increased when playing on hard surfaces owing to greater friction between the foot and the ground.<sup>11-17</sup> Clay courts seem to be a better option for ACL deficient players.

Limitations of this study include problems associated with questionnaires, and shortcomings related to retrospective determination of the preinjury performance level.

In summary, complete rupture of the ACL is a debilitating injury that causes significant alteration of knee joint kinematics. Untreated patients have joint instability, chronic articular degeneration, and knee dysfunction. Tennis specific limitations related to complete isolated ACL rupture were clearly identified. Tennis players with an ACL deficient knee showed significant impairment of subjective sport performance, limitation in landing after smashing, limitation in stopping and changing direction, limitation when playing a three set singles match, and limitation in playing on a hard court surface compared with healthy controls.

#### Authors' affiliations

**J Maquirriain**, National High Performance Training Centre (CeNARD), Argentine Tennis Association, Buenos Aires, Argentina

**P J Megey**, San Isidro Hospital, Buenos Aires

Competing interests: none declared

#### REFERENCES

- 1 **Hawkins RJ**, Misamore GW, Merritt TR. Follow-up of the acute non-operative isolated anterior cruciate ligament tear. *Am J Sports Med*, 1986;**14**:205-10.
- 2 **Noyes FR**, Moaar PA, Mathews DS, *et al*. The symptomatic anterior cruciate-deficient knee. Part 1. The long-term functional disability in athletically active individuals. *J Bone Joint Surg Am* 1983;**65A**:154-62.
- 3 **Kühne CA**, Zettl RP, Nast-Kolb D. [Tennis injuries: pattern an incidence in competitive and leisure tennis sports. ] *Sportvel Sportschad*, 2004;**18**:85-89.
- 4 **Vahey TN**, Meyer SF, Shelbourne KD, *et al*. MR imaging of anterior cruciate ligament injuries. *MRI Clin North Am* 1994;**2**:365-80.
- 5 **Tegner Y**, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop* 1985;**198**:43-9.
- 6 **Mont MA**, LaPorte DM, Mullick T, *et al*. Tennis after total hip arthroplasty. *Am J Sports Med* 1999;**27**:1-5.
- 7 **Castiglia VC**. Programa experimental de cálculos estadísticos VCC 001. Módulo para estimación de N muestrales ([www.institutodemetodologia.com](http://www.institutodemetodologia.com)).
- 8 **McNair PJ**, Marshall RN. Landing characteristics in subjects with normal and anterior cruciate ligament deficient knee joints. *Arch Phys Med Rehabil* 1994;**75**:584-9.
- 9 **Czierniecki JM**, Lippert F, Olerund JE. A biomechanical evaluation of tibiofemoral rotation in anterior cruciate deficient knees during walking and running. *Am J Sports Med* 1988;**16**:327-31.
- 10 **Steele JR**, Brown JM. Effects of chronic anterior cruciate ligament deficiency on muscle activation patterns during an abrupt deceleration task. *Clin Biomech* 1999;**14**:247-57.
- 11 **Besier TF**, Lloyd DG, Cochrane JL, *et al*. External loading of the knee joint during running and cutting maneuvers. *Med Sci Sports Exerc* 2001;**33**:1168-75.
- 12 **Butler DL**, Noyes FR, Grood ES. Ligamentous restraints to anterior-posterior drawer in the human knee: a biomechanical study. *J Bone Joint Surg Am* 1980;**62A**:259-70.
- 13 **Sallay PI**, Feagin JA. Basic science of the crucial ligaments. In: McGinty JB, Caspari RB, Jackson RW, *et al*. *Operative arthroscopy*, 2nd edition. Philadelphia: Lippincott-Raven Publishers, 1996.
- 14 **Schabus R**. Reconstructed anterior cruciate ligaments in tennis players. In: Krahl H, Pieper HG, Kibler WB, *et al*, editors. *Tennis: sports medicine and science*. Düsseldorf: Walter Rau Verlag, 1995.
- 15 **Besier TF**, Lloyd DG, Ackland TR, *et al*. Anticipatory effects on knee joint loading during running and cutting maneuvers. *Med Sci Sports Exerc* 2001;**33**:1176-81.
- 16 **Besier TF**, Lloyd DG, Ackland TR. Muscle activation strategies at the knee during running and cutting maneuvers. *Med Sci Sports Exerc* 2003;**35**:119-27.
- 17 **Nigg BM**, Segesser B. The influence of playing surface on the load on the locomotor system and on football and tennis injuries. *Sports Med* 1988;**5**:375-85.

#### BJSM presents a new feature: Online First

In an innovative move, BJSM is now publishing selected papers *Online First* within days of acceptance. These unedited articles are posted on the BJSM website (<http://bjsm.bmjournals.com>) weekly and are citable from the moment they are first posted; they are also deposited in PubMed. Every article will be published in print in its final, edited version when space in an issue becomes available. All versions will remain accessible via the website.

These articles can be accessed via the BJSM homepage or by using standard author and keyword searches on BJSM Online, Google and PubMed.

Sign up for BJSM announcements (<http://bjsm.bmjournals.com/cgi/alerts/etoc>) to be notified when new papers are published Online First.